

A PROOF OF THE LAWSON CONJECTURE FOR MINIMAL TORI EMBEDDED IN $\mathbb{S}^3$

FERNANDO A. A. PIMENTEL

ABSTRACT. A peculiarity of the geometry of the euclidean 3-sphere $\mathbb{S}^3$ is that it allows for the existence of compact without boundary minimally immersed surfaces. Despite a wealthy of examples of such surfaces, the only known tori minimally embedded in $\mathbb{S}^3$ are the ones congruent to the Clifford torus. In 1970 Lawson conjectured that the Clifford torus is, up to congruences, the only torus minimally embedded in $\mathbb{S}^3$. We prove here Lawson conjecture to be true. Two results are instrumental to this work, namely, a characterization of the Clifford torus in terms of its first eigenfunctions ([MR]) and the assumption of a “two-piece property” to these tori: every equator divides a torus minimally embedded in $\mathbb{S}^3$ in exactly two connected components ([R]).

1. INTRODUCTION

A special feature of the the euclidean 3-sphere $\mathbb{S}^3 = \{x \in \mathbb{R}^4 : |x| = 1\}$ on which it differs from the euclidean space $\mathbb{R}^3$ is that its topology does not obstruct the existence of compact without boundary minimally immersed surfaces. In fact, a plethora of examples of such surfaces are known to exist. Thus it was shown in [HL] by Hsiang-Lawson the existence of infinitely many immersed tori. On the other hand, examples of embedded minimal surfaces of arbitrary genus are constructed in [L1] by Lawson, with additional examples of such surfaces being supplemented by Karcher-Pinkall-Sterling in [KPS].

As regards embedded minimal surfaces, any impression of superabundance in examples suggested by the papers cited above evanesces as soon as one realizes that all minimally embedded n -tori hitherto known must have plenty of symmetries, that are inherited from their construction process. For instance, the only known minimally embedded torus is, up to congruences, the Clifford torus (see Def. 2.1), which has all the symmetries compatible with its topology.

Some results indicate that a minimally embedded torus must indeed be very symmetrical. Thus Lawson [L4] proved that any embedded minimal torus must be unknotted. In 1995, Ros [R] extended the restrictions on minimally embedded tori by proving that they have a special kind of *two-piece property* (see also Def. 3.1):

If M is an embedded compact minimal surface in $\mathbb{S}^3$, then every totally geodesic equator of $\mathbb{S}^3$ divides M in exactly two connected components.

This last result will be used extensively in this work.

It is well known that the euclidean coordinates $x_1, \dots, x_4$ of a minimal immersion M in $\mathbb{S}^3$ are eigenfunctions of the Laplacian on M (see section 5). With respect to this, we state a characterization of the Clifford torus due to Montiel-Ros [MR] that shall be useful to us as well:

The only minimal torus immersed into $\mathbb{S}^3$ by its first eigenfunctions is the Clifford torus.

This remarkable result relates the Yau conjecture, which asserts that any minimal embedding of a compact surface is by the first eigenfunctions and the Lawson conjecture:

CONJECTURE (Lawson, [L4]): *a torus minimally embedded in $\mathbb{S}^3$ is congruent to the Clifford torus.*

Statements of these conjectures may be found in [Y] (both conjectures) and [L4] (original statement of Lawson conjecture). Our aim here is to produce a proof of Lawson conjecture (Theorem 5.1).

Our approach to this problem is twofold, based on the results cited above in [R] and [MR]. Along Section 3 we use the *two-piece property* to classify the intersection of an equator $S(v)$ with a torus M with the *two-piece property* and strictly negative Gauss-Kronecker curvature (see Section 2 and Def. 3.1 for notation and basic definitions) in four types:

Proposition 3.1 *Let M be a $C^{2,\alpha}$ torus embedded in $\mathbb{S}^3$ that has strictly negative Gauss-Kronecker curvature. Then M has the two-piece property if and only if for every equator $S(v)$ the intersection $S(v) \cap M$ is classified among the four types listed below:*

- (1) *an embedded closed curve,*
- (2) *two disjoint embedded closed curves,*
- (3) *two embedded closed curves bounding disjoint discs in $S(v)$ with only one point in common,*
- (4) *two embedded closed curves bounding disjoint discs in $S(v)$ with exactly two points in common.*

where a curve in a intersection of type 1 is nullhomotopic in M and the pair of curves in intersections of types 2, 3 or 4 are homotopic in M to a same generator of $\pi_1(M)$ for any choice of the pair of curves in type 4 intersections (see Remark 3.3).

Diagrams of these intersections are sketched in Appendix A.

Then we show that for each torus M as above there exists an equator $S(v)$ such that the components of $M \setminus S(v)$ are two tubes with common border disjoint smooth closed curves in $S(v)$:

Lemma 3.4 *There exists an equator $S(v)$ such that $S(v) \cap M$ is of type 2.*

These results establish a kind of “topological symmetry” strong enough to provide us the geometric support needed in our next step: granted the existence of a minimal torus not congruent to the Clifford torus, construct a sequence of minimal tori converging in $C^{2,\alpha}$ to the Clifford torus. The *two-piece property* will be used to control the convergence process from which the sequence above shall be built (see Section 4). We then prove Lemma 4.5 and its corollary:

Corollary 4.1 *Let $0 < \alpha < 1$. If there exists a torus M minimally embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus T then there also exists a sequence (X_k) of $C^{2,\alpha}$ diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 such that each $M_k = X_k(T)$ is a torus minimally embedded in $\mathbb{S}^3$ noncongruent to the Clifford torus and $\lim_{k \rightarrow \infty} \|X_k - I\|_{C^{2,\alpha}} = 0$.*

Above it is mentioned the annulus $A_2 = \{x \in \mathbb{R}^4 \mid 1/2 < |x| < 2\}$ in connection with a canonical extension of maps of $\mathbb{S}^3$ to maps of A_2 , defined in Section 4. This seemingly extraneous apparatus is in fact pertinent. The extension of maps from the ambient space $\mathbb{S}^3$ to A_2 will somewhat simplify our approach due to the vector space structure of $\mathbb{R}^4$ thus incorporated. This will enable us to make use of ordinary Holder spaces of functions on regions in a euclidean space.

A crucial point in the proof of Lemma 4.5 and its Corollary 4.1 is to ensure that the tori M_k are not congruent to the Clifford torus. With respect to this, we remark the role of Lemma 3.6. It asserts that some closed curve in a type 2 intersection given by Lemma 3.4 stated above is not congruent to any closed curve in the Clifford torus T , assumed M minimal and noncongruent to T . This fact will be used to distinguish the tori M_k from the Clifford torus.

On the other hand, from the supracited characterization of the Clifford torus in [MR], it will be proven in Section 5 that the Clifford torus is isolated in $C^{2,\alpha}$. In order to do that, it will be shown that whenever (M_k) is a sequence of embedded tori noncongruent to the Clifford torus T converging to T in $C^{2,\alpha}$ then their first eigenvalues will converge to two, the first eigenvalue of T . But the first eigenfunctions of M_k will not converge to the first eigenfunctions of T as well. From these remarks a contradiction will arise and the result below will follow:

Lemma 5.1 *Let $0 < \alpha < 1$. Then there does not exist any sequence (X_k) of diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 such that each $M_k = X_k(T)$ is a torus minimally embedded in $\mathbb{S}^3$ noncongruent to the Clifford torus T and $\lim_{k \rightarrow \infty} \|X_k - I\|_{C^{2,\alpha}} = 0$.*

The main result in this paper (Theorem 5.1) is an immediate consequence of Corollary 4.1 and Lemma 5.1.

Part of this work was in the Doctoral thesis of the author at UFC, Brazil. I would like to thank A. Gervásio Colares for his orientation and continuous encouragement. I would also like to thank H. Rosenberg and A. Ros for some very valuable discussions. I am specially grateful to Abdénago A. de Barros that introduced me to the Lawson conjecture and to the results in [R] and [MR]. Finally, I would like to express my gratitude to the Université Paris XII and the Ministère de la Recherche, France, for the post-doc position that enabled me to discuss my issues with some of the afore mentioned professors.

2. PRELIMINARIES

The elements of spherical geometry that we will need are sketched here and found in full detail in [Bg].

Let $\mathbb{S}^3 \subset \mathbb{R}^4$ be the set of unit vectors of $\mathbb{R}^4$. A totally geodesic equator, hereafter called equator, is a totally geodesic big 2-sphere of $\mathbb{S}^3$. Let $\langle \cdot, \cdot \rangle$ be the inner product of $\mathbb{R}^4$. Following [R], we let correspond to any unit vector $v \in \mathbb{R}^4$ the equator $S(v) = \{p \in \mathbb{S}^3 \mid \langle v, p \rangle = 0\}$ and the half spheres $H_+(v) = \{p \in \mathbb{S}^3 \mid \langle v, p \rangle > 0\}$ and $H_-(v) = \{p \in \mathbb{S}^3 \mid \langle v, p \rangle < 0\}$.

For “distance between points” it is meant here the intrinsic distance of $\mathbb{S}^3$. A circle C in an equator $S(v)$ of $\mathbb{S}^3$ is a closed curve in $S(v)$ whose points equidist from a center, that is, a distinguished point in $S(v)$. According to this definition, the circle C has two centers p_1, p_2 in the equator $S(v)$, which are antipodal points (i.e., $p_1 = -p_2$) as it is easily seen. The curvature of C is given by $|\cotg(r)|$, where

r is the distance from C to any of its centers (see in [Sp], vol 3, the definition of curvature of a curve). If C has positive curvature then we can define the normal vector field along $C \subset S(v)$ as the unit vector field along C , normal to C and tangent to $S(v)$, that points towards the connected component of $S(v) \setminus C$ containing the center closer to C .

When a circle C has zero curvature it is called a geodesic or great circle. We can perform a rotation of an equator in $\mathbb{S}^3$ around any geodesic in it, so a geodesic is contained in infinitely many equators with a pair of centers in each equator.

Tori dealt with in this paper are $C^{2,\alpha}$ maps $X : S^1 \times S^1 \rightarrow \mathbb{S}^3$, immersions (when dX is injective) and embeddings (when it is further assumed that X is injective). Defined along these tori are the principal curvatures, denoted by k_1 and k_2 , the mean curvature $H = (k_1 + k_2)/2$ and the Gauss-Kronecker curvature $S = k_1 k_2$. We recall that the intrinsic or sectional curvature of surfaces in $\mathbb{S}^3$ is given by $K = 1 + S$. Hence the Clifford torus, with principal curvatures $k_1 = 1$ and $k_2 = -1$ for a suitably choice of a normal vector field is a minimal ($H \equiv 0$) flat ($K \equiv 0$) torus as we will verify below. We refer to [Sp], vols. 3 and 4, for the definitions and a throughout analysis of the curvature functions along immersed surfaces.

Definition 2.1. *We identify $\mathbb{R}^4$ with $\mathbb{C}^2$ in the usual manner. The Clifford torus is the compact surface in $\mathbb{S}^3$ given by*

$$\{(u, v) \in \mathbb{C}^2 : |u| = |v| = \sqrt{2}/2\}.$$

The orbits of the action of the groups $O^+(\mathbb{C}) \times \text{id}_{\mathbb{C}}$ and $\text{id}_{\mathbb{C}} \times O^+(\mathbb{C})$ in the Clifford torus(see [Bg], 18.8.6) are two families of circles parameterized by $u \rightarrow (u, v_0)$ and $v \rightarrow (u_0, v)$ for $|u| = |v| = |u_0| = |v_0| = \sqrt{2}/2$. These circles have curvature equal to 1 and circles in different families are orthogonal at their only common point. Moreover, a simmetry argument implies that a vector field normal to any of these circles is normal to the Clifford torus too. Thus these families of curves are the two systems of lines of curvature of the Clifford torus.

3. THE TWO-PIECE PROPERTY

Below it is presented a property concerning equators and subsets of $\mathbb{S}^3$ already mentioned in the Introduction.

Definition 3.1. *We say that a set $S \subset \mathbb{S}^3$ has the two-piece property when $S \setminus S(v)$ has exactly 2 connected components for each equator $S(v)$.*

The Clifford torus is itself an instance of surface with the *two-piece property*. In order to verify this either apply Ros' Theorem (see Introduction) or follow this straightforward reasoning: cyclides of Dupin (i.e., tori whose lines of curvature in the two systems are circles, e.g., any tori of revolution) are known to be divided in two connected components by any equator transversal to them at some point (see [B]). The Clifford torus is a cyclide of Dupin (see Section 2) transversal to any equator (since it has strictly negative Gauss-Kronecker curvature (see Sect. 2), it is not contained in any halfspace $H_+(v)$ or $H_-(v)$; or one could move the equator $S(v)$ until it touches the Clifford torus by one of its sides).

Nevertheless, being a cyclide of Dupin is so strong a constraint for a tori to have the *two-piece property*. Hereafter we will obtain weaker conditions that imply the *two-piece property* for tori in $\mathbb{S}^3$.

Firstly, we remark that tori with the *two-piece property* can not have points of positive Gauss-Kronecker curvature: consider an equator tangent to a torus at a point of positive Gauss-Kronecker curvature. A suitable motion of the equator shall increase the number of connected components of the torus bordered by the equator.

Besides, minimally immersed tori do not have umbilical points (i.e., where $k_1 = k_2 = 0$), see Lemma 1.4 and the ensuing comments in [L1]. So these tori have strictly negative Gauss-Kronecker curvature too. Consequently, we will restrict our analysis to tori with strictly negative Gauss-Kronecker curvature.

Now we observe that it can be proved that the properties of tori that concern us here, i.e., embeddedness, negativity of the Gauss-Kronecker curvature and the *two-piece property*, rest undisturbed after small perturbations. Thus

Lemma 3.1. *The set of embedded $C^{2,\alpha}$ tori with two-piece property and strictly negative Gauss-Kronecker curvature is open in the space of $C^{2,\alpha}$ tori endowed with the $C^{2,\alpha}$ topology.*

Remark 3.1. *A general definition of the $C^{2,\alpha}$ topology for maps from M to N , where M, N are manifolds, can be found in [H], section 2.1.*

The following Lemmas classify the curves in the intersection of equators and tori with the *two-piece property* and strictly negative Gauss-Kronecker curvature embedded in $\mathbb{S}^3$ in terms of their number, shape and homotopy class (we refer to [M] for the nomenclature and results assumed hereon).

The terminology “meridians” and “longitudes” is well established and refers to the two systems of canonical generators of the fundamental group $\pi_1(\overline{M})$ of a torus $\overline{M}$ embedded in $\mathbb{R}^3$. Informally, they are the curves that run once in the proper sense around the torus, thus linking any homotopically nontrivial curve in the bounded component of $\mathbb{R}^3 \setminus \overline{M}$ (meridians) or in its unbounded component (longitudes) (see [A], p.108).

Analogously, there exists a natural system of generators of the fundamental group $\pi_1(M)$ of a torus M in $\mathbb{S}^3$. They may be called meridians and longitudes as its counterparts on tori embedded in $\mathbb{R}^3$, after a choice of a normal vector field along M . We define them as the images of the lines of curvature of the Clifford torus T by some diffeomorphism X of $\mathbb{S}^3$ such that $X(T) = M$ that preserves the chosen orientation. *For the sake of simplicity we shall adopt “generator” for “canonical generator”.*

For a disc it is meant any subset homeomorphic to $\{x \in \mathbb{R}^2 \mid |x| < 1\}$ and a tube is a disc minus one point. *In what follows M is an embedded $C^{2,\alpha}$ torus, $0 < \alpha \leq 1$, with strictly negative Gauss-Kronecker curvature and the two-piece property.*

Lemma 3.2. *If $S(v)$ is an equator that is not tangent to M at any point then $S(v) \cap M$ either is an only embedded closed curve bounding a disc in M or is the union of two disjoint embedded closed curves homotopic in M to a same generator of $\pi_1(M)$.*

Proof. Obviously $S(v) \cap M$ is not empty. In fact, it is the union of disjoint closed embedded regular curves since M is an embedded compact smooth surface, and M is strictly transversal to $S(v)$.

Let M^+ , M^- be the closure of the connected components of $M \setminus S(v)$, with $M^+ \setminus S(v) \subset H^+(v)$ and $M^- \setminus S(v) \subset H^-(v)$. Thus, by the *two-piece property* and the embeddedness assumed for M , both M^+ and M^- are compact connected

orientable surfaces with boundary. It is well known that M^+ and M^- are then embedded orientable compact surfaces in $\mathbb{S}^3$ without a finite number of disjoint discs whose borders, i.e., $S(v) \cap M^+$ and $S(v) \cap M^-$, are the embedded closed curves in the intersection $S(v) \cap M$ (see prob. 4, p. 144, in [Sg]). We classify below $S(v) \cap M$ according to the topology of M^+ .

- (1) M^+ is a sphere without an only disc. Then $S(v) \cap M$ contains an only embedded closed curve bounding a disc in M , that is, $S(v) \cap M^+$;
- (2) M^+ is a sphere without exactly two disjoint discs. Then M^- must be itself an sphere without two discs, otherwise M is not a torus. Thus the intersection of M^+ with $S(v)$, i.e., $S(v) \cap M$, is the disjoint union of two curves homotopic to a same generator of $\pi_1(M)$;
- (3) M^+ is a sphere without $n \geq 3$ discs. Then at least M^- is an sphere without n discs and M is an n -torus, a contradiction;
- (4) M^+ is a torus without an only disc. In order that M be a torus, M^- must be a sphere without a disc. Thus item (4) is similar to item (1);
- (5) M^+ is a k -torus, $k \geq 1$ without $n \geq 2$ discs. Then M^- is at least an sphere without n discs and M is not a torus. A contradiction (see item (3)). $\square$

If M is tangent to an equator $S(v)$ then the points of tangency are isolated from each other (we remember that M has strictly negative Gauss-Kronecker curvature, so points of tangency p in $S(v) \cap M$ are nondegenerate saddle points regarding M as a graphic over $S(v)$ near p). So one can carry out a $C^{2,\alpha}$ deformation of M by perturbing it in a neighborhood of its points of tangency in order that $S(v)$ is not tangent to the deformed torus $\overline{M}$ anymore. If the deformation is sufficiently small, we can guarantee that $\overline{M}$ is embedded, has strictly negative Gauss-Kronecker curvature and the *two-piece property* (see Lemma 3.1).

Remark 3.2. *Here's a local description of the deformation above: in a small neighborhood of a point of tangency p the intersection $S(v) \cap M$ is the union of two curves crossing at p at a nonnull angle forming an "x" shape (nondegenerate saddle point). We denote by q_1, q_2, q_3, q_4 the points at the extremities of the "x". It may be supposed that the points q_i are in a circle in $S(v)$ with center p such that around the circle q_1 lies between q_2 and q_4 , q_2 lies between q_1 and q_3 , etc. Thus the "x" is formed by smooth arcs connecting q_1 to q_3 and q_2 to q_4 crossing at a nonnull angle at the point $p \neq q_i$. We assume the deformation near p restricted to a small ball B with center p such that $q_1, \dots, q_4 \notin B$. Since $\overline{M}$ is not tangent to $S(v)$, locally $S(v) \cap \overline{M}$ is the disjoint union of two curves that either connect q_1 to q_2 and q_3 to q_4 or connect q_1 to q_4 and q_2 to q_3 . See in [A], p. 101, a nice picture of this bifurcation.*

Lemma 3.3. *An equator $S(v)$ can be tangent to M at most at two points.*

Proof. Suppose $S(v)$ tangent to M . The assumptions on M imply that $S(v) \setminus M$ is the union of open regions whose borders are closed curves embedded in $S(v)$. We remark that these curves are not necessarily smooth, for points of tangency may belong to some of them.

Let $\mathcal{C}_1, \mathcal{C}_2$ be the connected components of $\mathbb{S}^3 \setminus M$. If both $\mathcal{C}_1 \cap S(v)$ and $\mathcal{C}_2 \cap S(v)$ have only one connected component, then $S(v) \cap M$ contains an only closed curve embedded in $S(v)$, that is to say, the border of these connected components in $S(v)$.

Besides, which is contrary to our assumptions, $S(v)$ is not tangent to M at any point because near points of tangency curves in $S(v) \cap M$ must have the “x” shape described in Remark 3.2.

Thus we can assume that $\mathcal{C}_1 \cap S(v)$ has at least two connected components. Firstly, we assume that $\mathcal{C}_1 \cap S(v)$ has exactly two connected components bordered, respectively, by the closed curves l_1 and l_2 embedded in $S(v)$ (so both components are discs in $S(v)$). Observe that $l_1 \cap l_2$ may have only points where $S(v)$ and M are tangent so it is a discrete (possibly even empty) set ($S(v)$ is tangent to M at a discrete set of points because M has strictly negative Gauss-Kronecker curvature). In fact, if $l_1 \cap l_2$ had a point q where $S(v)$ and M are strictly transversal then a line in $S(v)$ would cross l_1 and l_2 at q without interchanging connected components of $\mathbb{S}^3 \setminus M$, a contradiction since M is supposedly embedded.

Admit that $l_1 \cap l_2$ has three or more points $q_1, q_2, \dots, q_n$ disposed sequentially along l_1 (and thus along l_2 as can be seen). Thus the union of closed arcs in l_1 and l_2 with extremities q_1 and q_2 is a closed embedded curve λ_1 bounding a disc in $S(v)$. Analogously embedded closed curves λ_2 and λ_3 are obtained from arcs in l_1 and l_2 connecting q_2 to q_3 and q_3 to q_4 (or to q_1 , if there are only 3 points in $l_1 \cap l_2$), respectively.

Now we will deform M into an embedded $C^{2,\alpha}$ torus $\overline{M}$ with the *two-piece property* and strictly negative Gauss-Kronecker curvature such that $\overline{M}$ and $S(v)$ are not tangent (Lemma 3.1). As seen in Remark 3.2, along the deformation the curves can bifurcate at the tangency points so that λ_1 , λ_2 and λ_3 are taken to disjoint embedded closed curves $\overline{\lambda}_1$, $\overline{\lambda}_2$ and $\overline{\lambda}_3$, respectively. This contradicts Lemma 3.2.

Now let $\mathcal{C}_1 \cap S(v)$ have two or more connected components bordered by embedded closed curves $l_1, l_2, \dots, l_n$, $n \geq 3$, where as above distinct curves do not cross each other (they border connected components of $\mathcal{C}_1$) and may only have a discrete set of points in common. In order to finish the proof of this lemma we can either proceed as above or in a more straightforward approach separate the borders of these connected components by a small perturbation of M around the points of tangency between M and $S(v)$, which contradicts Lemma 3.2 again. $\square$

Lemmas above are synthetized in the following proposition:

Proposition 3.1. *Let M be a $C^{2,\alpha}$ torus embedded in $\mathbb{S}^3$ that has strictly negative Gauss-Kronecker curvature. Then M has the two-piece property if and only if for every equator $S(v)$ the intersection $S(v) \cap M$ is classified among the four types listed below:*

- (1) *an embedded closed curve,*
- (2) *two disjoint embedded closed curves,*
- (3) *two embedded closed curves bounding disjoint discs in $S(v)$ with only one point in common,*
- (4) *two embedded closed curves bounding disjoint discs in $S(v)$ with exactly two points in common.*

where a curve in a intersection of type 1 is nullhomotopic in M and the pair of curves in intersections of types 2, 3 or 4 are homotopic in M to a same generator of $\pi_1(M)$ for any choice of the pair of curves in type 4 intersections (see Remark 3.3).

Remark 3.3. *A type 4 intersection may be described as follows: $S(v) \cap M$ is the union of two embedded smooth closed curves C_1 and C_2 crossing at two points P*

and Q where these curves are strictly transversal. Then let l_1, l_2, l_3 and l_4 be arcs in $S(v)$ with common extremities P and Q such that, seen as pointsets, $l_1 \cup l_3 = C_1$, $l_2 \cup l_4 = C_2$ and $l_i \cap l_j = \{P, Q\}$ when $i \neq j$. Then one has two choices for the pair of embedded curves bounding disjoint discs in $S(v)$ meeting only at two points: the curves of trace $l_1 \cup l_2$ and $l_3 \cup l_4$; or the curves of trace $l_1 \cup l_4$ and $l_2 \cup l_3$.

Proof of Prop. 3.1. If the intersection $S(v) \cap M$ is of the types above then necessarily $S(v)$ divides M in two connected components. Indeed, the constraints on the shape and homotopy class of the curves in the intersections completely characterize the topology of $H_+(v) \cap M$ and $H_-(v) \cap M$ up to a reflection with respect to $S(v)$ (see appendix A). Thus $M \setminus S(v)$ either is a disjoint union of a disc with a torus without a disc (type 1), or a disjoint union of two tubes (type 2), and so on.

For the reciprocal, that was mostly the object of Lemmas 3.2 and 3.3, we assume that M has the *two-piece property*. What is left to prove concerns the homotopy class of curves of types 3 and 4. But this characterization follows from the proof of Lemma 3.3. In fact, it was proved there that if $S(v)$ is tangent to M , then $S(v) \cap M$ is the union of two embedded closed curves l_1 and l_2 with at least one and at most two points in common bounding disjoint discs in $S(v)$. Now deform M into an embedded torus $\overline{M}$ with the *two-piece property* and strictly negative Gauss-Kronecker curvature. From Remark 3.2 and the proof of Lemma 3.3, we may assume that l_1 and l_2 are taken continuously along the deformation into disjoint closed curves $\bar{l}_1$ and $\bar{l}_2$. By Lemma 3.2 these curves are homotopic generators of $\pi_1(\overline{M})$, so l_1 and l_2 are homotopic generators too. $\square$

Remark 3.4. *Types 1, 2, 3 or 4 also describe the possible intersections of an ordinary torus of revolution in $\mathbb{R}^3$ with a plane to which it is transversal at some point. This fact can be verified with the aid of the sketches provided in appendix A.*

We now assume that $S(v)$ is tangent to an embedded torus M with strictly negative Gauss-Kronecker curvature and the *two-piece property*. Let p be one of the points where M and $S(v)$ are tangent. In a neighborhood of p the intersection $S(v) \cap M$ contains two disjoint curves crossing p at a nonnull angle (since M has strictly negative Gauss-Kronecker curvature). This angle is formed by the asymptotic directions of M at p (see [Sp]). Admit that a geodesic γ passes through p without pointing to an asymptotic direction. Near p , a rotation of $S(v)$ around γ splits the curves in $S(v) \cap M$ into two connected smooth curves with γ between them, regardless the sense of the rotation. While one of these curves does not touches γ near p the other one must be tangent to γ at p (which passes through p depends on the sense of the rotation).

If γ does not pass through the tangency point p then the bifurcation depends on the sense of the rotation and proceeds exactly as described in Remark 3.2. Indeed, Remark 3.2 treats a particular case in which p is a center of the geodesic γ , i.e., a point that equidists from the geodesic. Thus, in the notation of Remark 3.2, if q_1, q_2, q_3 and q_4 are the extremities of the “x” shape in $S(v) \cap M$ with center p , then we may assume that a small rotation of $S(v)$ around γ to an equator $S(\bar{v})$ takes q_1, q_2, q_3 and q_4 to points $\bar{q}_1, \bar{q}_2, \bar{q}_3$ and $\bar{q}_4$, respectively, in $S(\bar{v}) \cap M$, such that in a neighborhood of p , according to the sense of the rotation, either $\bar{q}_1$ is connected to $\bar{q}_2$ and $\bar{q}_3$ to $\bar{q}_4$ by disjoint arcs in $S(\bar{v}) \cap M$, or $\bar{q}_1$ is connected to $\bar{q}_4$ and $\bar{q}_2$ to $\bar{q}_3$.

Thus let $\gamma \subset S(v)$ be a geodesic that does not point to asymptotic directions of M at points where M is tangent to $S(v)$. If $S(\bar{v})$ is obtained by a small rotation of

$S(v)$ around γ then $S(\bar{v})$ is not tangent to γ , provided the rotation is sufficiently small. Then, as we have seen, near the points of tangency the intersection $S(\bar{v}) \cap M$ is the union of regular disjoint curves and outside a neighborhood of the points of tangency both M and $S(v)$ (hence also M and $S(\bar{v})$) are strictly transversal if the rotation is sufficiently small.

These remarks will be used in the proof of the following result:

Lemma 3.4. *There exists an equator $S(v)$ such that $S(v) \cap M$ is of type 2.*

Proof. Firstly, we assume that M is not invariant by the antipodal map. Thus let p be a point in M such that $-p$ does not belong to M . Let $S(\bar{v})$ the equator tangent to M at p .

If $S(\bar{v}) \cap M$ is of type 3, let D_1 and D_2 be the two disjoint discs in $S(\bar{v})$ bounded by closed curves in $S(\bar{v}) \cap M$ whose common point is p . Then assume that the geodesic $\gamma \subset S(\bar{v})$ passes through p without pointing to an asymptotic direction and does not cross D_1 and D_2 . Thus, near p , the geodesic γ lies between D_1 and D_2 . Hence, in a neighborhood of p , the intersection $S(v) \cap \gamma$ is the union of four arcs, namely l_1 , l_2 , L_1 and L_2 , with only common point the extremity p such that l_1 and l_2 are in the border of D_1 and L_1 , L_2 border D_2 (see Remark 3.2). Let $l_3 = \partial D_1 \setminus (l_1 \cup l_2)$ and $L_3 = \partial D_2 \setminus (L_1 \cup L_2)$.

A small rotation of $S(\bar{v})$ around γ into an equator $S(v)$ takes $(l_1 \cup l_2)$ and $L_1 \cup L_2$ into smooth curves l_α , L_α in $S(v) \setminus \gamma$, respectively. As seen above, these curves are smooth, disjoint and rest in different sides of γ . As regards the disjoint smooth curves l_3 and L_3 , these curves are taken by the small rotation into disjoint smooth curves in $S(\bar{v}) \cap M$, say l_β and L_β . We may assume that the extremities of l_α and L_α coincide with the extremities of l_β and L_β , respectively. Then provided the rotation is sufficiently small, $l_\alpha \cup l_\beta$ and $L_\alpha \cup L_\beta$ are the trace of disjoint embedded closed curves and $S(v) \cap M$ is of type 2.

If $S(\bar{v}) \cap M$ is of type 4 with $S(\bar{v})$ tangent to M at p and q then $S(\bar{v}) \setminus M$ contains among its components two disjoint discs D_1 and D_2 whose boundaries in $S(\bar{v}) \cap M$ are two closed curves with common points only p and q . As above let $\gamma \subset S(\bar{v})$ be a geodesic through p that does not point to an asymptotic direction and also separates D_1 from D_2 near p . Since M is not invariant by the antipodal map we may assume that $q \neq -p$ so we may also suppose that γ does not pass through q . We define l_1 , l_2 and L_1 , L_2 in a neighborhood of p as in the preceding argument for type 3 intersections. Analogously we denote by l_3 , l_4 , L_3 , $L_4 \subset S(\bar{v}) \cap M$ the arcs in a small neighborhood of q with a extremity in q such that $l_3 \cup l_4 \subset \partial D_1$ and $L_3 \cup L_4 \subset \partial D_2$. The sets $S_1 = \partial D_1 \setminus (l_1 \cup l_2 \cup l_3 \cup l_4)$ and $S_2 = \partial D_2 \setminus (L_1 \cup L_2 \cup L_3 \cup L_4)$ are the disjoint union of smooth arcs l_5 , $l_6 \subset \partial D_1$ and L_5 , $L_6 \subset \partial D_2$, respectively.

After a small rotation of $S(\bar{v})$ into $S(v)$ the curves l_1 , l_2 , L_1 , L_2 are taken to smooth disjoint curves l_α and L_α of $S(v) \cup M$ as in the previous case. We may choose the sense of the rotation (see discussion above and Remark 3.2) in order that l_3 , l_4 and L_3 , L_4 are taken, respectively, to smooth disjoint arcs l_β (obtained from l_3 , l_4) and L_β (obtained from L_3 , L_4) as described in Remark 3.2. Arcs l_5 , l_6 , L_5 and L_6 , are taken to smooth disjoint arcs l_μ , l_ν , L_μ and L_ν , respectively. So we can assume that $l_\alpha \cup l_\beta \cup l_\mu \cup l_\nu$ and $L_\alpha \cup L_\beta \cup L_\mu \cup L_\nu$ are the trace of disjoint closed curves in $S(v) \cap M$ that is thus of type 2.

Finally, let M be invariant by the antipodal map and $S(\bar{v})$ be tangent to M at the point p . Then $S(\bar{v})$ is tangent to M at the point $-p$ too. Thus $S(\bar{v})$ is of type 4 and there exists disjoint discs D_1 and D_2 that are components of $S(\bar{v}) \setminus M$

whose borders have common points p and $-p$. Let γ be a geodesic of $S(\bar{v})$ such that at p γ does not point to an asymptotic direction and separates D_1 from D_2 in a neighborhood of p . One may verify that the antipodal map preserves connected components of $\mathbb{S}^3 \setminus M$. Thus the antipodal invariance of M implies that at $-p$ the geodesic γ does not point to an asymptotic direction and locally separates D_1 from D_2 . As in the analysis of a type 3 intersection, once a small rotation of $S(\bar{v})$ into $S(v)$ is performed, the discs D_1 and D_2 are set apart into connected components of $S(v) \setminus M$ with disjoint boundaries. So $S(v) \cap M$ is of type 2. $\square$

Lemma 3.5. *Let T be the Clifford torus and $S(v)$ an equator tangent to T . Let $(v_n) \subset \mathbb{S}^3$ be a sequence in $\mathbb{S}^3$ such that $v_n \rightarrow v$ and each $S(v_n)$ is not tangent to T at any point. Then*

- (1) *each intersection $S(v_n) \cap T$ is of type two;*
- (2) *if λ_n is a closed curve in $S(v_n) \cap T$ then there exists points $p_n, q_n \in \lambda_n$ such that $k_n(p_n) \rightarrow \infty$ and $k_n(q_n) \rightarrow 0$ when $n \rightarrow \infty$, where k_n is the curvature of λ_n .*

Proof.

(1) $\mathbb{S}^3 \setminus T$ is the union of two connected components that are invariant by the antipodal map. Now assume that $S(v_n) \cap T$ is of type 1. The connected components S_1, S_2 of $S(v_n) \setminus T$ are in different connected components of $\mathbb{S}^3 \setminus T$. Moreover, the antipodal map takes S_1 to S_2 and vice versa. A contradiction.

(2) In the proof of Lemma 3.4 it was argued that the invariance of T by the antipodal map implies that $S(v) \cap T$ is of type 4. Besides, $S(v) \cap T$ has all the compatible symmetries. A curve in $S(v) \cap T$ separating two regions must bend equally into the two regions, so its curvature must be equal to zero. The intersection $S(v) \cap T$ is then the union of two geodesics crossing at the antipodal points $p, -p$ at an angle equal to $\pi/2$. By taking a subsequence, if necessary, we may assume that the sequence (λ_n) converges to arcs of geodesics in $S(v) \cap T$ bounding a component of $S(v) \setminus T$. One may choose points $q_n \in \lambda_n$ away from $p, -p$ at least a sufficiently small $\epsilon > 0$. Thus $k_n(q_n) \rightarrow 0$. Assume that the curvature of λ_n remains bounded when $n \rightarrow \infty$. Thus a subsequence of (λ_n) converges to a curve λ that at least belongs to $C^{1,\alpha}$ for $0 \leq \alpha < 1$ (since $C^2(S^1)$ is compact in $C^{1,\alpha}(S^1)$, see Theorem 1.31 in [Ad]). A contradiction, because (λ_n) converges to arcs of geodesic that meet at an angle equal to $\pi/2$. So there exists the sequence (p_n) defined above. $\square$

Lemma 3.6. *Let M be a minimal torus embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus T . Then there exists an equator $S(v_0)$ such that $S(v_0) \cap M$ is of type 2 and contains a closed curve that is not congruent to any closed curve in a intersection $S(v) \cap T$ of type 2.*

Remark 3.5. *Lemma 3.6 seems quite straightforward since M and T are noncongruent real analytic tori (see Lemma 1.1 in [L1]). Its proof below would be simpler had we used a fact communicated to us by H. Rosenberg, namely, that whenever a torus minimally embedded in $\mathbb{S}^3$ contains a geodesic it is congruent to the Clifford torus.*

Proof of Lemma 3.6. Firstly, we recall that the torus M has strictly negative Gauss-Kronecker curvature and the *two-piece property* (by Lemma 1.4 in [L1] and Theorem 2 in [R]; see also Int.). We also observe that there exists an equator $S(w)$ such that $S(w) \cap M$ is of type 2 (Lemma 3.4). Suppose that whenever $S(w) \cap M$

is of type 2 then both closed curves in $S(w) \cap M$ are congruent to closed curves in intersections $S(\bar{w}_1) \cap T$ and $S(\bar{w}_2) \cap T$, respectively.

We may choose $v_n \in \mathbb{S}^3$ such that $S(v_n) \cap M$ is of type 2 for every n . Moreover, we can also assume that $v_n \rightarrow v$, and that $S(v)$ is tangent to M (just take $v_0 \in \mathbb{S}^3$ such that $S(v_0) \cap M$ is of type 2 and move $S(v_0)$ until it becomes tangent to M).

Thus $S(v_n)$ converges to $S(v)$ and curves in $S(v) \cap M$ must be the limit (in the sense discussed in item 2 of the proof of Lemma 3.5) of closed curves congruent to curves in intersections $S(w) \cap T$. Besides curves in $S(v) \cap M$ are not in C^2 because of their singularities at the tangency points. Thus $S(v) \cap M$ *must be congruent to an intersection $S(\bar{v}) \cap T$ of type 4 such that $S(v) \cap M$ is the intersection of two geodesics of $\mathbb{S}^3$ crossing at an angle equal to $\pi/2$* (see proof of Lemma 3.5).

We consider then the set S of points p where M is tangent to equators $S(v_p)$ such that $S(v_p) \cap M$ is the union of two geodesics crossing at an angle equal to $\pi/2$. This set is readily seen to be closed and not empty, by Lemma 3.5 and the remarks above. Let $p \in S$ and V be a small neighborhood of p in M . Let $S(v_q)$ be an equator tangent to M at some point q in V . We have the following situations to analyze:

(1) If $S(v_q) \cap M$ is of type 3 then we proceed as in the proof of Lemma 3.4. We rotate $S(v_q)$ around a geodesic $\gamma \subset S(v_q)$ through q that lets the embedded closed curves in $S(v_q) \cap M$ in the closures of different connected components of $S(v_q) \setminus \gamma$. As was seen above the rotated equators have intersections of type 2 with M whenever the rotation is sufficiently small. Our hypothesis thus implies that $S(v_q) \cap M$ is the union of two geodesics meeting at two points at an angle equal to $\pi/2$, a contradiction.

(2) Now suppose that $S(v_q) \cap M$ is of type 4. If $S(v_p)$ is the equator tangent to M in p then there exists a geodesic γ in $S(v_p)$ such that the closure of each component of $S(v_p) \setminus \gamma$ contains an embedded closed curve. Hence when V is sufficiently small and $q \in V$ the same happens to any intersection $S(v_q) \cap M$ of type 4. Thus we rotate the equator $S(v_q)$ around a geodesic in corresponding to γ above in order to obtain type 2 intersections as in the previous case. Then we proceed as in item 1 above.

So S is open and closed in M besides being not empty. Hence $S = M$ and the asymptotic curves of M (i.e., curves whose normal curvature is equal to zero, see [Sp], vol. 3) are geodesics of $\mathbb{S}^3$. Then M is an embedded flat minimal torus (quick proof: if $p \in M$, consider a sequence R_n of rectangles with vertex p whose sides are arcs of asymptotic curves; then assume that the length of the sides of R_n tends to zero when $n \rightarrow \infty$ and apply the Gauss-Bonnet formula). According to a well known result due in [CCK] and [L3], the minimal torus M must be the Clifford torus, a contradiction. $\square$

The next result is quite straightforward as well and may also be obtained from the characterization of the Clifford torus in Remark 3.5.

Lemma 3.7. *Let M be a torus minimally embedded in $\mathbb{S}^3$ and assume that $S(v) \cap M$ is of type 2. Then none of the closed curves in $S(v) \cap M$ is a geodesic of $\mathbb{S}^3$.*

Proof. Let us admit that $\lambda_1 \subset S(v) \cap M$ is a geodesic of $\mathbb{S}^3$. We perform a rotation of $S(v)$ around λ_1 until an equator $S(w)$ obtained from the rotation of $S(v)$ becomes tangent to M at a point p (such equator does exist because $\lambda_1 \subset M$). If $p \notin \lambda_1$ then $S(w) \cap M$ is not of types 1, 2, 3 or 4 and M does not have the *two-piece property*

(see Prop. 3.1), which contradicts Ros' Theorem (i.e., Theorem 1.2 in [R]). Thus $p \in \lambda_1$.

Now we observe that $S(v) \cap M$ is of type 2. Hence $S(v) \setminus \lambda_1$ is the union of two discs D_1 and D_2 and we may assume from the definition of type 2 intersections that $D_1 \cap M = \emptyset$ and that $D_2 \cap M$ is a closed curve. Along the rotation, until the first point of tangency, the intersections of the equators with M remain of type 2. Hence the disc rotated from D_2 contains a closed curve and the disc rotated from D_1 has empty intersection with M . Let the discs D_1, D_2 be taken by the rotation above to discs $D_1^*, D_2^* \subset S(w)$, respectively, which are the components of $S(w) \setminus \lambda_1$. Thus, $D_1^* \cap M = \emptyset$ (or M does not have strictly negative Gauss-Kronecker curvature) and, consequently, the point of tangency p cannot be a nondegenerate saddle point (a curve through p in $S(w) \cap M$ must be in the closure of D_2^*). This contradicts the already cited fact (for example, in the proof of the preceding lemma) that M has strictly negative Gauss-Kronecker curvature. $\square$

4. SEQUENCES OF MINIMAL TORI CONVERGING TO THE CLIFFORD TORUS

Let the open annulus $A_2 = \{x \in \mathbb{R}^4 : 1/2 < |x| < 2\}$ and $F : A_2 \rightarrow A_2$ be a $C^{2,\alpha}$ map. We let $D^j = \partial/\partial x_j$, $D^{jk} = \partial^2/\partial x_j \partial x_k$ and set

$$[F]_\alpha = \sup_{\substack{x, y \in A_2 \\ x \neq y}} \frac{|F(x) - F(y)|}{|x - y|^\alpha}, \quad (1)$$

$$\begin{aligned} \|F\|_{C^{2,\alpha}} &= \sup_{A_2} |F(x)| + \max_{1 \leq j \leq 4} \sup_{A_2} |D^j F(x)| + \max_{1 \leq j, k \leq 4} \sup_{A_2} |D^{jk} F(x)| \\ &\quad + \max_{1 \leq j, k \leq 4} [D^{jk} F]_\alpha, \end{aligned} \quad (2)$$

a seminorm and the Holder $C^{2,\alpha}$ norm in A_2 , respectively, where $|\cdot|$ is the euclidean norm in $\mathbb{R}^4$.

Remark 4.1. *If F can be extended in C^2 to a map G defined in the closure A_2° then it is easily seen that $\|F\|_{C^{2,\alpha}}$ and $\|G\|_{C^{2,\alpha}(A_2^\circ)}$ coincide.*

The purpose of the following lemma is to serve as a compactness criterion for what follows. Let X, Y , where $X \subset Y$, be topological spaces. We observe that only in Lemma 4.1 below the term “embedding” refers to the identity map $I : X \rightarrow Y$ when I is continuous. If there also holds that $I(X)$ is precompact in Y then we say that X is compactly embedded in Y .

Any $C^{2,\alpha}$ diffeomorphism ξ of $\mathbb{S}^3$ is canonically extended to a $C^{2,\alpha}$ diffeomorphism X of A_2 : if $v \in \mathbb{S}^3$ and $1/2 < r < 2$ let $X(rv) = r\xi(v)$. Thus

Lemma 4.1. *Let $\mathcal{X}^\alpha$ be the set of $C^{2,\alpha}$ diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 . If $0 < \nu < \lambda \leq 1$ then the embedding $\mathcal{X}^\lambda \rightarrow \mathcal{X}^\nu$ exists and is compact.*

Proof. Lemma 4.1 follows directly from Remark 4.1 and Theorem 1.31 in [Ad]. $\square$

We let now $I : A_2 \rightarrow A_2$ be the identity map in A_2 and define the following functional on the diffeomorphisms of A_2 :

$$\tau^\alpha(X) = \|X - I\|_{C^{2,\alpha}} + \|X^{-1} - I\|_{C^{2,\alpha}}, \quad (3)$$

where $0 < \alpha \leq 1$.

Lemma 4.2. *If $0 < \alpha \leq 1$ let (X_k) be a sequence of $C^{2,\alpha}$ diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 as above. Assume that $\tau^\alpha(X_k) \leq C < \infty \forall k \in \mathbb{N}$. Let $0 < \beta < \alpha$. Then a subsequence of (X_k) converges in $C^{2,\beta}$ to a diffeomorphism X of $\mathbb{S}^3$ canonically extended to A_2 . Moreover, besides being a map in $C^{2,\beta}$, X is also a $C^{2,\alpha}$ diffeomorphism and $\tau^\alpha(X) \leq C$. Lastly, if $X_k \rightarrow X$ in $C^{2,\alpha}$, then X is a $C^{2,\alpha}$ diffeomorphism of $\mathbb{S}^3$ canonically extended to A_2 , and there exists $\lim_{k \rightarrow \infty} \tau^\alpha(X_k) = \tau^\alpha(X)$.*

Proof. Since the norms $\|X_k - I\|_{C^{2,\alpha}}$ are bounded for all k , the sequence (X_k) is bounded in $C^{2,\alpha}$. By Lemma 4.1 there exists a subsequence, denoted (X_{k_l}) , that converges in $C^{2,\beta}$ to a map $X : A_2 \rightarrow A_2$. Now we consider the sequence $(X_{k_l}^{-1})$ obtained from the subsequence above. From the fact that the norms $\|X_{k_l}^{-1} - I\|_{C^{2,\alpha}}$ are also bounded for all k and Lemma 4.1 we have that a subsequence of $(X_{k_l}^{-1})$, denoted $(X_{k_l}^{-1})$ as well, converges in $C^{2,\beta}$ to a $C^{2,\beta}$ map $X^* : A_2 \rightarrow A_2$. A continuity argument implies that exists $X^{-1} = X^*$. Hence X is a $C^{2,\beta}$ diffeomorphism of $\mathbb{S}^3$ canonically extended to A_2 .

In fact, X is a $C^{2,\alpha}$ diffeomorphism. By a routine argument, let $1 \leq i, j \leq 4$: since $\lim_{l \rightarrow \infty} D^{ij} X_{k_l}(x) = D^{ij} X(x)$ then for every pair $x, y \in A_2$, $x \neq y$, there holds that

$$\liminf_{l \rightarrow \infty} [D^{ij}(X_{k_l} - I)]_\alpha \geq \lim_{l \rightarrow \infty} \frac{|D^{ij}(X_{k_l} - I)(x) - D^{ij}(X_{k_l} - I)(y)|}{|x - y|^\alpha} = \frac{|D^{ij}(X - I)(x) - D^{ij}(X - I)(y)|}{|x - y|^\alpha}.$$

Thus both $[D^{ij}(X - I)]_\alpha$, $\|X\|_{C^{2,\alpha}} < \infty$. In fact, $\liminf_{l \rightarrow \infty} [D^{ij}(X_{k_l} - I)]_\alpha \geq [D^{ij}(X - I)]_\alpha$. Besides, since it was assumed that $X_{k_l} \rightarrow X$ in $C^{2,\beta}$ for some $0 < \beta < \alpha$, the other terms in the expression of $\|X_{k_l} - I\|_{C^{2,\alpha}}$ converge to their counterparts in the expression of $\|X - I\|_{C^{2,\alpha}}$. From these remarks there follows that $\liminf_{l \rightarrow \infty} \|X_{k_l} - I\|_{C^{2,\alpha}} \geq \|X - I\|_{C^{2,\alpha}}$. Analogously we have the corresponding inequality for the inverse map: $\liminf_{l \rightarrow \infty} \|X_{k_l}^{-1} - I\|_{C^{2,\alpha}} \geq \|X^{-1} - I\|_{C^{2,\alpha}}$. Thus $C \geq \liminf_{l \rightarrow \infty} \tau^\alpha(X_{k_l}) \geq \tau^\alpha(X)$.

If $X_k \rightarrow X$ in $C^{2,\alpha}$ then $(X_k - I)$ and $X_k^{-1} - I$ converge to $X - I$ and $X^{-1} - I$ in $C^{2,\alpha}$, respectively, since $\tau^\alpha(X_k)$ is bounded, and the Lemma follows. $\square$

The Clifford torus T (see Def. 2.1) is reticulated by the longitudes and meridians below:

$$\phi_n^k(\theta) = \frac{\sqrt{2}}{2}(\cos(\frac{\pi k}{n}), \sin(\frac{\pi k}{n}), \cos(\theta), \sin(\theta)) \quad (4)$$

$$\psi_n^k(\theta) = \frac{\sqrt{2}}{2}(\cos(\theta), \sin(\theta), \cos(\frac{\pi k}{n}), \sin(\frac{\pi k}{n})) \quad (5)$$

where $\theta \in [0, 2\pi)$ and $0 \leq k < 2n$. The components of T minus the longitudes and meridians above are the $4n^2$ squares in T with vertices the points

$$p_n^{jk} = \frac{\sqrt{2}}{2}(\cos(\frac{\pi j}{n}), \sin(\frac{\pi j}{n}), \cos(\frac{\pi k}{n}), \sin(\frac{\pi k}{n})). \quad (6)$$

Now let $v_0 = (0, 1, 0, 0)$. Thus $S(v_0) \cap T$ is of type 2 and is the disjoint union of the lines of curvature ϕ_n^0 and ϕ_n^n .

Hereafter we assume the curves in $S(v_0)$ parameterized by arclength. So let λ be a $C^{2,1}$ closed curve embedded in the equator $S(v_0)$ and Ω_n^λ be the set of $C^{2,1}$

diffeomorphisms X of $\mathbb{S}^3$ canonically extended to A_2 that let $S(v_0)$ invariant such that both $X(\phi_n^0)$ is congruent to λ (seen as a pointset) and the image $X(T)$ of the Clifford torus has mean curvature equal to zero at the points $X(p_n^{jk})$ (abusing of the notation, here T , λ , ϕ_n^0 , etc., when arguments of maps, must be regarded as pointsets).

Remark 4.2. *We observe that Ω_n^λ is not empty. In fact, it is seen that there exists a diffeomorphism ξ of $S(v_0)$ taking ϕ_n^0 to λ . Since the set of the p_n^{jk} is discrete, we may assume, by perturbing the diffeomorphism ξ , if necessary, that $\xi(T)$ has mean curvature zero at the points $\xi(p_n^{jk})$. This diffeomorphism is then extended canonically to a diffeomorphism X of A_2 as described above.*

We now define some functions on the set of closed $C^{2,1}$ curves λ embedded in $S(v_0)$. Firstly, we set

$$\delta_n(\lambda) = \inf\{\tau^1(X) : X \in \Omega_n^\lambda\}. \quad (7)$$

Thus $\delta_n(\lambda) = 0$ if and only if λ is congruent to ϕ_n^0 .

If $0 < \alpha < 1$ and M is an unknotted $C^{2,\alpha}$ torus (see [L4]) embedded in $\mathbb{S}^3$ let Λ_α^M be the set of $C^{2,\alpha}$ diffeomorphisms Y of $\mathbb{S}^3$ canonically extended to A_2 such that $Y(T)$ is congruent to M . Let now $\Omega_{n,\alpha}^{\lambda,\mathcal{K}}$ be the closure in $C^{2,\alpha}$ of the set of diffeomorphisms $X \in \Omega_n^\lambda$ such that $\tau^1(X) < \mathcal{K}$. We remark that $\Omega_{n,\alpha}^{\lambda,\mathcal{K}} \subset \Omega_n^\lambda$ by Lemma 4.2. Let $0 < \alpha < 1$ and $\mathcal{K} > \delta_n(\lambda)$, and set

$$\xi_{n,\alpha}^\mathcal{K}(\lambda) = \inf\{\tau^\alpha(Y) : Y \in \Lambda_\alpha^{X(T)} \text{ for some } X \in \Omega_{n,\alpha}^{\lambda,\mathcal{K}}\}. \quad (8)$$

Given an arclength parameterized arc μ with trace in $S(v_0)$ we obtain a closed arclength parameterized curve λ in $S(v_0)$ by joining the extremities of μ by an arclength parameterized arc ν in $S(v_0)$ and letting λ be the arclength parameterized curve whose trace is the union of the traces of μ and ν . We may suppose that λ is in $C^{2,1}$ by assuming that both $\mu, \nu \in C^{2,1}$ and that μ, ν join regularly (i.e., $C^{2,1}$) at their extremities. We denote this construction by the product $\lambda = \mu\nu$, where it is transparent that we identify distinct arclength parameterizations of λ .

Let λ be a closed C^3 curve embedded in $S(v_0)$ parameterized by arclength and $\mathcal{K} \geq \delta_n(\lambda) + 1$. Since λ is closed, we may consider it parameterized for all s by letting $\lambda(s) = \lambda(s \bmod 2L)$, where $2L$ is its length. We set $\lambda_t = \lambda|_{[-(1-t)L, (1-t)L]}$, $0 < t < 1$, for the restriction of λ to the interval $[-(1-t)L, (1-t)L]$. Let $\mathcal{S}_{n,\alpha}^t$, $0 < \alpha, t < 1$, be the set of $C^{2,1}$ arcs μ with the same extremities as λ_t such that the product $\lambda_t\mu$ is a closed $C^{2,1}$ curve embedded in $S(v_0)$ and $\xi_{n,\alpha}^\mathcal{K}(\lambda_t\mu)$ is defined. This set is not empty because it contains the arc of λ whose trace is $\lambda \setminus \lambda_t$. We set

$$\Phi_{n,\alpha}^\lambda(t) = \inf_{\mu \in \mathcal{S}_{n,\alpha}^t} \xi_{n,\alpha}^\mathcal{K}(\lambda_t\mu) \quad (9)$$

if $t \in (0, 1)$ and let $\Phi_{n,\alpha}^\lambda(0) = \xi_{n,\alpha}^\mathcal{K}(\lambda)$.

Lemma 4.3. *The function $\Phi_{n,\alpha}^\lambda$ is continuous and nonincreasing.*

Proof. Let $t_2 > t_1$ and $\mu \in \mathcal{S}_{n,\alpha}^{t_1}$. Thus $(\lambda_{t_1}\mu) \setminus \lambda_{t_2} \in \mathcal{S}_{n,\alpha}^{t_2}$ and, consequently, $\Phi_{n,\alpha}^\lambda$ is nonincreasing. Let $(\mu_k^t) \subset \mathcal{S}_{n,\alpha}^t$ be a minimizing sequence for $\Phi_{n,\alpha}^\lambda(t)$, i.e., $\lim_{k \rightarrow \infty} \xi_{n,\alpha}^\mathcal{K}(\lambda_t\mu_k^t) = \Phi_{n,\alpha}^\lambda(t)$. Taking into account that $\Phi_{n,\alpha}^\lambda$ is nonincreasing we admit that $\lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t) > \Phi_{n,\alpha}^\lambda(t_0)$ and let $2\epsilon = \lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t) - \Phi_{n,\alpha}^\lambda(t_0)$. If k is sufficiently large then $\xi_{n,\alpha}^\mathcal{K}(\lambda_{t_0}\mu_k^{t_0}) - \Phi_{n,\alpha}^\lambda(t_0) < \epsilon$. Thus there exists $(X, Y) \in$

$\Omega_{n,\alpha}^{\lambda_{t_0}\mu_k^{t_0},\mathcal{K}} \times \Lambda_\alpha^{X(T)}$ such that $\tau^\alpha(Y) - \Phi_{n,\alpha}^\lambda(t_0) < \epsilon$. Since $\Omega_{n,\alpha}^{\lambda_{t_0}\mu_k^{t_0},\mathcal{K}}$ is the closure in $C^{2,\alpha}$ of $C^{2,1}$ maps $Z \in \Omega_n^{\lambda_{t_0}\mu_k^{t_0}}$ such that $\tau^1(Z) < \mathcal{K}$, then we may assume that $\mathcal{K} > \tau^1(X)$. Now we recall that λ is a C^3 curve: thus $\lambda_{t_0-\delta} \rightarrow \lambda_{t_0}$ in $C^{2,1}$ (see Remark 3.1) and we may admit for each δ the existence of $\mu_\delta \in \mathcal{S}_{n,\alpha}^{t_0-\delta}$ such that $\lambda_{t_0-\delta}\mu_\delta \rightarrow \lambda_{t_0}\mu_k^{t_0}$ in $C^{2,1}$ when $\delta \rightarrow 0$ and thus also in $C^{2,\alpha}$. A continuity argument implies that if δ is sufficiently small then there are diffeomorphisms $(X_\delta, Y_\delta) \in \Omega_{n,\alpha}^{\lambda_{t_0-\delta}\mu_\delta,\mathcal{K}} \times \Lambda_\alpha^{X_\delta(T)}$ (since we may choose X_δ in order that $\tau^1(X_\delta) < \mathcal{K}$) near (X, Y) in $C^{2,1} \times C^{2,\alpha}$ such that $\tau^\alpha(Y_\delta) - \Phi_{n,\alpha}^\lambda(t_0) < \epsilon$, hence $\lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t) - \tau^\alpha(Y_\delta) > \epsilon$. On the other hand, $\tau^\alpha(Y_\delta) \geq \xi_{n,\alpha}^\mathcal{K}(\lambda_{t_0-\delta}\mu_\delta) \geq \Phi_{n,\alpha}^\lambda(t_0 - \delta) \geq \lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t)$, a contradiction. Thus $\lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t) = \Phi_{n,\alpha}^\lambda(t_0)$. *Mutatis mutandis*, a slight modification of the argument above proves that $\lim_{t \rightarrow t_0} \Phi_{n,\alpha}^\lambda(t) = \Phi_{n,\alpha}^\lambda(t_0)$ even when $t_0 = 0$. $\square$

Let M be a minimal torus embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus T . We may assume that $S(v_0) \cap M$ is of type 2 (by Lemma 3.4 associated with a congruence of $\mathbb{S}^3$). Since $S(v_0) \cap M$ is real analytic so are the curves in $S(v_0) \cap M$. Thus, by Lemma 3.6, we may assume that there exists a closed real analytic curve $\lambda^M \subset S(v_0) \cap M$ such that λ^M is not congruent to any closed curve in an intersection $S(v) \cap T$. Moreover, we also know that λ^M is not a geodesic of $\mathbb{S}^3$ (see Lemma 3.7).

Let X_M be a $C^{2,1}$ diffeomorphism of $\mathbb{S}^3$ extended to A_2 that lets $S(v_0)$ invariant such that $X_M(\phi_n^0)$ is congruent to λ^M and $X_M(T)$ is congruent to M . Such a diffeomorphism does exist, since M is unknotted (cf. [L4]) and λ^M is a $C^{2,1}$ embedded canonical generator of $\pi_1(M)$. Thus $X_M \in \Omega_n^{\lambda^M}$ and $\delta_n(\lambda^M) \leq \tau^1(X_M)$ for all $n \in \mathbb{N}$.

We choose an arclength parameterization of λ^M in order that the curvature of λ^M assumes its maximum $k_{\max}$ (which is necessarily positive since λ^M is not a geodesic) at $\lambda^M(0)$. As regards the Clifford torus T , there exists $v \in \mathbb{S}^3$ such that a real analytic embedded closed curve $\mu^T \subset S(v) \cap T$ has a critical point of curvature $k_{\max}$ (Lemma 3.5) at $\mu^T(0)$ for a suitably chosen arclength parameterization of μ^T . Thus there also exists a diffeomorphism X_T such that $X_T \in \Omega_n^{\mu^T}$ for all n since we may choose $X_T(T)$ congruent to T . Hence $\xi_{n,\alpha}^\mathcal{K}(\mu^T) = 0 \forall n \in \mathbb{N}$ and $\mathcal{K} > 0$.

For lemmas 4.4 and 4.5 we let $0 < \alpha < 1$ as usual and set

$$\mathcal{K} = \max(\tau^1(X_M), \tau^1(X_T)) + 1,$$

thus $\mathcal{K} > \delta_n(\lambda^M)$ as required.

Lemma 4.4. *Let M be a minimal torus embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus and λ^M be the closed curve defined above. Then there exists a constant $C > 0$ such that $\Phi_{n,\alpha}^{\lambda^M}(0) \geq C \forall n$. Besides, regardless n , $\lim_{t \rightarrow 1} \Phi_{n,\alpha}^{\lambda^M}(t) = 0$.*

Proof. Suppose that $\inf_{n \in \mathbb{N}} \Phi_{n,\alpha}^{\lambda^M}(0) = 0$. Then either (1) there exists a subsequence (n_k) such that $\lim_{k \rightarrow \infty} \Phi_{n_k,\alpha}^{\lambda^M}(0) = 0$ or (2) $\Phi_{n_0,\alpha}^{\lambda^M}(0) = 0$ for some $n_0 \in \mathbb{N}$. If only (2) happens just consider $n > n_0$ in the hypothesis of the lemma, though the reasoning below also applies to this case.

Thus we may admit that (1) occurs and then take maps $(X_{n_k}, Y_{n_k}) \in \Omega_{n_k,\alpha}^{\lambda^M,\mathcal{K}} \times \Lambda_\alpha^{X_{n_k}}$ such that $0 \leq \tau^\alpha(Y_{n_k}) - \Phi_{n_k,\alpha}^{\lambda^M}(0) < 1/k$. Thus by Lemma 4.2 there

holds that for $0 < \beta < \alpha$ subsequences of (X_{n_k}) and (Y_{n_k}) converge in $C^{2,\alpha}$ and $C^{2,\beta}$ to diffeomorphisms $X, Y \in C^{2,1} \times C^{2,\alpha}$, respectively. By Lemma 4.2, $0 = \lim_{k \rightarrow \infty} \tau^\alpha(Y_{n_k}) \geq \tau^\alpha(Y) \geq 0$. Thus Y is the identity map. Since $X_{n_k}(T)$ is congruent to $Y_{n_k}(T)$, there holds that $X(T)$ is congruent to the Clifford torus. Moreover, $S(v_0) \cap X(T)$ contains a closed curve congruent to λ^M , which contradicts the definition of λ^M .

We will now prove that $\lim_{t \rightarrow 1} \Phi_{n,\alpha}^{\lambda^M}(t) = 0$. Let λ_t^M, μ_t^T be arcs of the curves λ^M, μ^T , respectively, obtained from λ^M, μ^T as described in the proof of Lemma 4.3. In order to prove that $\lim_{t \rightarrow 1} \Phi_{n,\alpha}^{\lambda^M}(t) = 0 \forall n$ we verify the existence of arcs ν_t such that the product $\lambda_t^M \nu_t \rightarrow \mu^T$ in $C^{2,1}$ when $t \rightarrow 1$. Indeed, both arcs λ_t^M, μ_t^T are C^3 curves with curvature equal to $k_{\max}$ at $\lambda_t^M(0), \mu_t^T(0)$, respectively, which are also critical points of the curvature function along their respective curves. Thus, up to congruences, the distance from λ_t^M to μ_t^T in the $C^{2,1}$ space of those arcs of curves (that is known to be metrizable, see [H], chapter 1) tends to zero when $t \rightarrow 1$. So for t sufficiently close to 1 we can join the extremities of λ_t^M with an arc ν_t near $\mu_t^T \setminus \mu_t^T$ in $C^{2,1}$ such that $\nu_t \rightarrow \mu^T \setminus \mu^T(0)$ in $C^{2,1}$ when $t \rightarrow 1$, and the convergence $\lambda_t^M \nu_t \rightarrow \mu^T$ in $C^{2,1}$ follows. Hence, there exists diffeomorphisms X_t converging to X_T in $C^{2,1}$ when $t \rightarrow 1$ such that $X_t(T) \cap S(v_0)$ contains a closed curve congruent to $\lambda_t^M \nu_t$ (just perturb X_T in a neighborhood of ϕ_n^0). Consequently, $\lim_{t \rightarrow 1} \tau^1(X_t) = \tau^1(X_T) < \mathcal{K}$. Now we observe that $\xi_{n,\alpha}^{\mathcal{K}}(\mu^T) = 0$ (since $X_T(T)$ is congruent to the Clifford torus so the identity map $I \in \Lambda_\alpha^{X(T)}$; besides, $\tau^1(X_T) < \mathcal{K}$ from the definition of $\mathcal{K}$). Hence we can assume the existence of a $C^{2,\alpha}$ diffeomorphism Y_t of $\mathbb{S}^3$ canonically extended to A_2 associated with X_t that converges to the identity map I in $C^{2,\alpha}$ when $t \rightarrow 1$ such that $Y_t(T)$ is congruent to $X_t(T)$ (e.g., one can take $Y_t = X_t \circ X_T^{-1}$). Thus from lemmas 4.2 and 4.3 there holds that

$$0 \leq \lim_{t \rightarrow 1} \Phi_{n,\alpha}^{\lambda^M}(t) \leq \lim_{t \rightarrow 1} \xi_{n,\alpha}^{\mathcal{K}}(\lambda_t^M \nu_t) \leq \lim_{t \rightarrow 1} \tau^\alpha(Y_t) = \tau^\alpha(I) = 0,$$

and the lemma is proved. $\square$

We then extend continuously $\Phi_{n,\alpha}^{\lambda^M}$ to the interval $[0, 1]$ by letting $\Phi_{n,\alpha}^{\lambda^M}(1) = 0$.

Lemma 4.5. *Let M be a minimal torus embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus T and C be the constant provided by Lemma 4.4. Then for every $\rho \in [0, C]$ there exists a torus M_ρ minimally embedded in $\mathbb{S}^3$ that is congruent to T only when $\rho = 0$, and a $C^{2,\alpha}$ diffeomorphism Y_ρ of $\mathbb{S}^3$ canonically extended to A_2 satisfying $\tau^\alpha(Y_\rho) = \rho$ such that $Y_\rho(T)$ is congruent to M_ρ .*

Proof. Let λ^M be the closed curve embedded in $\mathbb{S}^3$ defined above. By Lemma 4.4 and the definition of the function $\Phi_{n,\alpha}^{\lambda^M}$ there holds that $\Phi_{n,\alpha}^{\lambda^M}(t_n^\rho) = \rho$ for some $t_n^\rho \in [0, 1]$. So for every $n \in \mathbb{N}$ and $\rho \in [0, C]$ there exists diffeomorphisms $(X_{n,\rho}, Y_{n,\rho}) \in \Omega_{n,\alpha}^{C_\rho^\rho, \mathcal{K}} \times \Lambda_\alpha^{X_{n,\rho}(T)}$, where $C_n^\rho = \lambda_{t_n^\rho}^M \mu_{t_n^\rho}^T$, such that $0 \leq \tau^\alpha(Y_{n,\rho}) - \rho < 1/n$.

Let $0 < \beta < \alpha$. We may assume that subsequences $(X_{n_k,\rho})$, and $(Y_{n_k,\rho})$ of the sequences above converge in $C^{2,\alpha}$ and $C^{2,\beta}$, respectively, to diffeomorphisms $X_\rho \in C^{2,1}$, $Y_\rho \in C^{2,\alpha}$ such that $\tau^1(X_\rho) \leq \mathcal{K}$, $\tau^\alpha(Y_\rho) \leq \rho$ (cf. Lemma 4.2) and $X_\rho(T)$ is congruent to $Y_\rho(T)$. Indeed, $\tau^\alpha(Y_\rho) = \rho$. In order to prove this, we will suppose that $\tau^\alpha(Y_\rho) < \rho$. Since $X_{n_k,\rho} \rightarrow X_\rho$ in $C^{2,\alpha}$ we may assume that there exists diffeomorphisms $Y_{n_k,\rho}^* \in \Lambda_\alpha^{X_{n_k,\rho}(T)}$ converging to Y_ρ in $C^{2,\alpha}$ (for instance, take $Y_{n_k,\rho}^* = X_{n_k,\rho} \circ X_\rho^{-1} \circ Y_\rho$). Hence, by Lemma 4.2 and the definition of $\Lambda_\alpha^{X_{n_k,\rho}(T)}$, there holds that $\rho \leq \lim_{k \rightarrow \infty} \tau^\alpha(Y_{n_k,\rho}^*) = \tau^\alpha(Y_\rho) < \rho$, a contradiction.

Let $M_\rho = Y_\rho(T)$. Then M_ρ is a $C^{2,1}$ torus embedded in $\mathbb{S}^3$. Identifying $X_{n,\rho}(T)$ and $Y_{n,\rho}(T)$ (that are congruent) there holds that $Y_{n,\rho}(T)$ has mean curvature equal to zero at the points $X_{n,\rho}(p_n^{jk})$, $1 \leq j, k \leq n$. We remark that the points $X_\rho(p_n^{jk})$, $n \in \mathbb{N}$, are dense in M_ρ . Thus convergence in $C^{2,\alpha}$ implies that M_ρ is minimal.

Suppose that M_ρ is congruent to the Clifford torus T for some $\rho \neq 0$. Then $I(T)$ is congruent to $X_\rho(T)$. As above there exists $C^{2,\alpha}$ diffeomorphisms $Y_{n_k,\rho}^*$ converging to I in $C^{2,\alpha}$ such that $Y_{n_k,\rho}^*(T) = X_{n_k,\rho}(T)$, i.e., $Y_{n_k,\rho}^* \in \Lambda_\alpha^{X_{n_k,\rho}(T)}$. A contradiction because $\lim_{k \rightarrow \infty} \tau^\alpha(Y_{n_k,\rho}^*) = 0 < \rho$. $\square$

The following corollary follows immediately from Lemma 4.5:

Corollary 4.1. *Let $0 < \alpha < 1$. If there exists a torus M minimally embedded in $\mathbb{S}^3$ that is not congruent to the Clifford torus T then there also exists a sequence (X_k) of $C^{2,\alpha}$ diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 such that each $M_k = X_k(T)$ is a torus minimally embedded in $\mathbb{S}^3$ noncongruent to the Clifford torus and $\lim_{k \rightarrow \infty} \|X_k - I\|_{C^{2,\alpha}} = 0$.*

5. MAIN RESULT

Corollary 4.1 implies the main result in this work if it is further proved that the Clifford torus is isolated in $C^{2,\alpha}$. In order to accomplish this we shall deal with the classical Hilbert spaces $L^2(M)$ and $H^1(M)$ of functions along a riemannian manifold $(M; \langle \cdot, \cdot \rangle)$. They are obtained by the completion of $C(M)$ and $C^1(M)$ with respect to the norms derived from the respective inner products:

$$\langle f, g \rangle_{L^2(M)} = \int_M f g \, dM, \quad (10)$$

$$\langle f, g \rangle_{H^1(M)} = \langle f, g \rangle_{L^2(M)} + \int_M \langle \text{grad}_M f, \text{grad}_M g \rangle \, dM. \quad (11)$$

We refer to [Be] for the properties of these spaces that we will need below, as those regarding the eigenvalues of the Laplace-Beltrami operator Δ_M on C^2 functions along $M^2 \subset \mathbb{S}^3$, i.e., the nonnegative λ for which the equation $\Delta_M u + \lambda u = 0$ has a nontrivial solution $u \in C^\infty(M)$ (u is then called an eigenfunction of λ). We recall the variational characterization of the first positive eigenvalue:

$$\lambda_1(M) = \inf_{\substack{f \in C^1(M) \\ \int_M f \, dM = 0}} \frac{\int_M |\text{grad}_M f|^2 \, dM}{\int_M f^2 \, dM}. \quad (12)$$

Finally, we remark that M^2 is minimally immersed in $\mathbb{S}^3$ if and only if the coordinate functions of the immersion in the ambient euclidean space $\mathbb{R}^4$ are eigenfunctions of the Laplace-Beltrami operator in the induced metric with eigenvalue equal to 2 (cf. [L2], parag. 4, prop. 1).

Lemma 5.1. *Let $0 < \alpha < 1$. Then there does not exist any sequence (X_k) of diffeomorphisms of $\mathbb{S}^3$ canonically extended to A_2 such that each $M_k = X_k(T)$ is a torus minimally embedded in $\mathbb{S}^3$ noncongruent to the Clifford torus T and $\lim_{k \rightarrow \infty} \|X_k - I\|_{C^{2,\alpha}} = 0$.*

Proof. We admit the existence of the sequences (M_k) , (X_k) described above. Let $\lambda_1(M_k)$ be the first positive eigenvalue of the Laplace-Beltrami operator Δ_{M_k} .

From the lower bound for $\lambda_1(M_k)$ in [CW] and the remark above on the coordinate functions of minimal immersions in $\mathbb{S}^3$ there holds that $1 < \lambda_1(M_k) \leq 2$.

By taking a subsequence if necessary we may assume that $(\lambda_1(M_k))$ is a convergent sequence. It is clear that $1 \leq \lim_{k \rightarrow \infty} (\lambda_1(M_k)) \leq 2$. Let $f_k : M_k \rightarrow \mathbb{R}^4$ be a nontrivial eigenfunction of the first eigenvalue $\lambda_1(M_k)$ of M_k and let $g_k = f_k \circ X_k$. Assume that $\int_{M_k} f_k^2 dM_k = 1$. These assumptions imply that $\lambda_1(M_k) = \int_{M_k} |\text{grad}_{M_k} f_k|^2 dM_k$ (see eq. 12). Let $\lambda = \lim_{k \rightarrow \infty} \lambda(M_k)$. Given $\delta > 0$ the following inequality holds for $k > 0$ sufficiently large

$$0 \leq \langle g_k, g_k \rangle_{H^1(T)}^2 \leq 1 + \lambda + \delta, \quad (13)$$

since $\lambda_1(M_k) \rightarrow \lambda$ and the norm of the jacobian of X_k tends to 1 when $k \rightarrow \infty$. Hence the sequence (g_k) is bounded in $H^1(T)$. Thus we may admit by taking a subsequence if necessary that the sequence (g_k) converges weakly to a function g in $H^1(T)$. It is well known that the inclusion $H^1(T) \subset L^2(T)$ is continuous and compact (see [Be]). So the sequence (g_k) is pre-compact in $L^2(T)$ and hence we can find a subsequence (g_{k_i}) that converges weakly to g in $H^1(T)$ and strongly to a function $\bar{g}$ in $L^2(T)$. Thus g_{k_i} converges weakly to $\bar{g}$ in $L^2(T)$ too since strong convergence implies weak convergence. The continuity of the inclusion $H^1(T) \subset L^2(T)$ implies that $g = \bar{g}$. From standard regularization procedures and the assumptions on f_k , M_k and g_k , we obtain that g is a smooth function satisfying $\int_T g^2 dM_T = 1$ and $\Delta_T g + \lambda g = 0$. Thus $1 \leq \lambda \leq 2$ is an eigenvalue of the Clifford torus, that is known to have the least positive eigenvalue equal to 2 (cf. [MR]). Hence $\lambda = 2$.


Let $x = (x_1, x_2, x_3, x_4)$ be the coordinate functions of the Clifford torus T . We define analogously the coordinate functions $x^k = (x_1^k, x_2^k, x_3^k, x_4^k)$ of the minimal embedding $M_k \subset \mathbb{S}^3$. As was observed above, $\Delta_T x + 2x = \Delta_{M_k} x^k + 2x^k = 0$. The functions x_i are a basis for the eigenspace of the first eigenvalue of the Clifford torus (see [C]). Thus the function g defined in the previous paragraph, an eigenfunction of the first positive eigenvalue of the Clifford torus as was shown, is given by $g = \sum_{i=1}^4 a_i x_i$. Let $h_k = \sum_{i=1}^4 a_i x_i^k$. We now observe that $\lambda_1(M_k) < 2$ or M_k is the Clifford torus (Theorem 4 in [MR]). Hence if f_k is the nontrivial eigenfunction of the first eigenvalue $\lambda(M_k)$ described above then $\int_{M_k} f_k h_k dM_k = 0$. A continuity argument implies that $\int_T g^2 dM_T = 0$, a contradiction. $\square$

Theorem 5.1 follows from Cor. 4.5 and Lemma 5.1.


Theorem 5.1. *If M is a torus minimally embedded in $\mathbb{S}^3$ then M is congruent to the Clifford torus.*

APPENDIX A. INTERSECTIONS OF EQUATORS AND TORI IN $\mathbb{S}^3$


We sketch here all the possible intersections of an equator $S(v)$ in $\mathbb{S}^3$ and an embedded torus M with the *two-piece property*. For a better understanding of these pictures one may assume that we have performed a stereographical projection from $\mathbb{S}^3$ to $\mathbb{R}^3$ with north pole a point in $S(v) \setminus M$. Thus the equator $S(v)$ is taken to a plane and the image of M is an ordinary torus of $\mathbb{R}^3$. For types 3 and 4 we have assumed that the north pole of the stereographical projection is chosen in such a way that the curves in $S(v) \cap M$ are taken to meridians of the projected torus.




type 1



type 2



type 3



type 4

REFERENCES

- [A] C. C. Adams, *The Knot Book*, W. H. Freeman and Company, New York, 2000.
- [Ad] R. A. Adams, *Sobolev Spaces*, Academic Press, 1975.
- [B] T. F. Banchoff, The Spherical Two-Piece-Property and Tight Surfaces in Spheres, *J. Diff. Geom.*, **4** (1970), 193-205.
- [Be] P. H. Berard, *Spectral Geometry: Direct and Inverse Problems (Lect. Notes in Math., Vol. 1207)*, Springer, 1986.
- [Bg] M. Berger, *Geometry*, Springer, 1987, 2 vols.
- [CCK] S. S. Chern, M. P. do Carmo & S. Kobayashi, Minimal submanifolds of a sphere with second fundamental form of constant length, *Funct. Anal. and Related Fields*, (1970), 57-75.
- [C] I. Chavel, *Eigenvalues in Riemannian Geometry*, Academic Press, Inc. 1984.
- [CW] H. I. Choi & A. N. Wang, A first eigenvalue estimate for minimal hypersurfaces, *J. Diff. Geom.*, **18** (1983), 559-562.
- [H] M. W. Hirsch, *Differential Topology*, Springer 1976.
- [HL] W.-Y. Hsiang & H. B. Lawson, Jr., Minimal manifolds of low cohomogeneity, *J. Diff. Geom.* **5** (1971), 1-38.
- [KPS] H. Karcher, U. Pinkall & I. Sterling, New minimal surfaces in $\mathbb{S}^3$, *J. Diff. Geom.* **28** (1988), 169-185.
- [L1] H. B. Lawson, Jr., Complete minimal surfaces in $\mathbb{S}^3$, *Ann. of Math.* **92** (1970), 335-374.
- [L2] ———, *Lectures on Minimal Submanifolds*, Publish or Perish 1980.
- [L3] ———, Local rigidity theorem for minimal hypersurfaces, *Ann. of Math.* **89** (1969), 187-197.
- [L4] ———, The unknottedness of minimal embeddings, *Invent. Math.* **11** (1970), 183-187.
- [M] J. P. May, *A Concise Course in Algebraic Topology*, Univ. of Chic. Press 1999.
- [MR] S. Montiel & A. Ros, Minimal immersions of surfaces by the first eigenfunctions and conformal area, *Invent. Math.* **83** (1986), 153-166.
- [R] A. Ros, A two-piece property for compact minimal surfaces in a three sphere, *Indiana Univ. Math. J.* **44** (1995), no. 3, 841-849.
- [Sg] G. Springer, *Riemann Surfaces*, 2nd ed., Chelsea.
- [Sp] M. Spivak, *A comprehensive introduction to differential geometry*, Publish or Perish 1974, 5 vols.
- [Y] S.-T. Yau, Problem section. Seminar on Differential Geometry, *Ann. Math. Stud.*, vol. 102, Princeton Univ. Press 1982.

Fernando A. A. Pimentel: Departamento de Matemática, UFC, Campus do Pici
bloco 914, 60455-760 Fortaleza-Ce, Brazil pimentelf@gmail.com